

Canonical Phase Diagrams of the 1-D Falicov-Kimball Model at $T=0$

Z. Gajek

Institute of Low Temperature
and Structure Research of Polish Academy of Sciences,
P. O. Box 937, 50–950 Wrocław, Poland

J. Jędrzejewski

Institute of Theoretical Physics,
University of Wrocław,
Max Born Sq. 9, 50–204 Wrocław, Poland

*R. Lemański**

Institute of Low Temperature
and Structure Research of Polish Academy of Sciences,
P. O. Box 937, 50–950 Wrocław, Poland

Abstract

The Falicov-Kimball model of spinless quantum electrons hopping on a 1-dimensional lattice and of immobile classical ions occupying some lattice sites, with only intrasite coupling between those particles, have been studied at zero temperature by means of well-controlled numerical procedures. For selected values of the unique coupling parameter U the restricted phase diagrams (based on all the periodic configurations of localized particles (ions) with period not greater than 16 lattice constants, typically) have been constructed in the grand-canonical ensemble. Then these diagrams have been translated into the canonical ensemble. Compared to the diagrams obtained in other studies our ones contain more details, in particular they give better insight into the way the mixtures of periodic phases are formed. Our study has revealed several families of new characteristic phases like the generalized most homogeneous and the generalized crenel phases, a first example of a structural phase transition and a tendency to build up an additional symmetry – the hole-particle symmetry with respect to the ions (electrons) only, as U decreases.

*formerly known as R. Łyżwa

1 The model and the method

The work reported in this paper has been inspired by the recent results due to Freericks and Falicov [1], Farkašovský and Bat'ko [2], and Gruber et al. [3]. It is also a continuation of our earlier work [4]. All the papers mentioned are concerned with various properties of the ground state phase diagram of the two-state spinless Falicov-Kimball model. This is a lattice model of itinerant quantum electrons and immobile classical ions that are coupled via a simple on site interaction. It is a simplified version of the model that was put forward in 1969 by Falicov and Kimball [5] to describe conductor-insulator transitions. In the one-dimensional case, considered in this paper, the Hamiltonian corresponding to the model has the form

$$H_{\Lambda} = \sum_{x \in \Lambda} \left(-a_x^* a_{x+1} - a_{x+1}^* a_x - U w(x) a_x^* a_x \right), \quad (1)$$

where Λ is a finite piece of the one-dimensional lattice with periodic boundary conditions imposed, a_x^*, a_x are the creation and annihilation operators of itinerant spinless electrons and $w(x) = 0$ or 1 is the occupation number of the ions at the lattice site x . The function $\Lambda \ni x \rightarrow w(x)$, is called the *ion configuration*. Observe that the electrons can hop only between the nearest neighbour sites and the energy units have been chosen in such a way that the hopping rate is unity. Therefore in H_{Λ} there is only one independent parameter expressed in these units: the electron-ion coupling U . The total number of electrons $N_e = \sum_{x \in \Lambda} a_x^* a_x$ and the local occupation numbers $w(x)$ of the ions (hence the total number $N_i(w) = \sum_{x \in \Lambda} w(x)$ of the ions in the configuration w) are the conserved quantities.

The Falicov–Kimball model can be related to the celebrated Hubbard model [6], [7]. It has however many interpretations of its own [1], [6], [8], interesting enough to motivate intensive investigations of this model. In one of the interpretations proposed by Khomskii [8], Kennedy and Lieb [6] and adopted here it is a model of crystalline formation. As proved by Kennedy and Lieb [6] and by Brandt and Schmidt [9] the effective interaction between the ions, induced by the direct interaction with mobile electrons, leads to the

formation of periodic structures of the ions (superlattices). Another interpretation that is sound from the point of view of solid state physics, is in terms of mobile s -electrons and localized f -electrons in rare-earth compounds. This makes the model relevant for understanding mixed-valence phenomena [5], [8].

For a given ion configuration w the Hamiltonian (1) is the second quantized form of the *one-particle operator* $h(w)$ whose matrix elements are

$$[h(w)]_{xy} = -t_{xy} - Uw(x)\delta_{xy}, \quad (2)$$

where

$$t_{xy} = \begin{cases} 1 & \text{if } y = x \pm 1 \\ 0 & \text{otherwise.} \end{cases} \quad (3)$$

In the sequel by the *spectrum of the configuration* w we mean the spectrum of $h(w)$.

The states of the whole system are characterized by their ion configurations and the electrons' wave functions. If the ion configuration w and the total number of electrons N_e are specified, then in the ground state the electrons' wave function corresponds to the so called *Fermi sea*. The energy of this state, $E^U(w; N_e)$, called the *Fermi sea energy*, is equal simply to the sum of all the N_e lowest eigenvalues of $h(w)$. Such states of the system are referred to as *phases*. Suppose that we are given the total number N_e of the electrons and the total number N_i of the ions. In the ground state subject to these conditions the ions are arranged in a specific configuration $\tilde{w}$, the *ground state configuration*, and the electrons form the corresponding Fermi sea. This state of the whole system is referred to as the (N_e, N_i) -phase. Its energy, $E(N_e, N_i)$, called the *ground state energy*, is equal to $E^U(\tilde{w}; N_e)$. The largest eigenvalue of $h(\tilde{w})$ that contributes to $E^U(\tilde{w}; N_e)$, referred to as the *Fermi level*, is separated from the higher and lower parts of the spectrum of $\tilde{w}$ by the gaps $\Delta_u(\tilde{w})$ and $\Delta_l(\tilde{w})$, respectively. A phase that is the (N_e, N_i) -phase for some pair (N_e, N_i) is said to appear in the canonical phase diagram or is referred to as *stable phase*.

Our results refer exclusively to the *infinite system*. The thermodynamic limit is taken in such a way that the electron number per site $N_e/|\Lambda|$, the ion number per site $N_i/|\Lambda|$, the Fermi sea energy per site $E^U(w; N_e)/|\Lambda|$ and the ground state energy per

site $E(N_e, N_i)/|\Lambda|$ converge to the electron density ρ_e , the ion density ρ_i , the Fermi sea energy density $e^U(w; \rho_e)$ and the ground state energy density $e(\rho_e, \rho_i)$, respectively. Suppose that the ion configuration of the (ρ_e, ρ_i) -phase is $\tilde{w}$. If in the thermodynamic limit $\Delta_u(\tilde{w}) - \Delta_l(\tilde{w}) > 0$, then the (ρ_e, ρ_i) -phase is called the *insulator*, otherwise it is termed the *conductor*.

The possibility of studying directly infinite systems stems from the fact that for the periodic ion configuration w the Green function of the tight binding Schrödinger equation and consequently the density of states $n(w)$, ρ_e and $e^U(w; \rho_e)$ can be determined exactly. However the corresponding formulae depend on the band edges of the spectrum of w , which are zeroes of the polynomial whose order coincides with the period of w [10]. First these zeros have to be determined numerically and after that the quantities of interest, like the energy density $e^U(w; \rho_e)$, can be obtained by a numerical integration.

The hole-particle unitary transformations applied to the Hamiltonian (1) reveal the corresponding symmetries of the Fermi sea energy density $e^U(w; \rho_e)$. Thus, performing the hole-particle transformation with respect to the ions we obtain

$$e^{(U)}(w^*; \rho_e) = e^{(-U)}(w; \rho_e) - U\rho_e, \quad (4)$$

where w^* is the *adjoint* ion configuration of w , $w^*(x) = 1 - w(x)$, while with respect to the electrons we get

$$e^{(U)}(w; \rho_e) = e^{(-U)}(w; 1 - \rho_e) - U\rho_i(w). \quad (5)$$

Therefore we can reduce the range of ρ_e values considered to the interval $[0, 1/2]$ and we can also fix arbitrarily the sign of U . We choose $U > 0$, i. e. according to (1) the ions and electrons attract each other at the same lattice site. Thus it is natural to refer to the system with $\rho_e = \rho_i$ as *neutral*.

In this paper we study the properties of the Falicov-Kimball model at zero temperature by means of an approximate method, the *method of restricted phase diagrams*. It is the set of the states available for the system considered that is approximated (by a restricted

set of states), not the interactions. We concentrate mainly on the *ion subsystem*, that is, we try to determine how the ions are arranged in the ground state. This sort of approach has already been used, first in [11], then in [1], [2], [3], [4],

The method of restricted phase diagrams is straightforward in principle. In the *canonical ensemble*, we are interested in here, some specified values are assigned to the densities ρ_e and ρ_i and the task amounts to minimizing the set of functions $\{e(w, \rho_e) : w \in \mathcal{W}_{\rho_i}\}$ over the set $\mathcal{W}_{\rho_i}$ of *admissible configurations* that satisfy the condition $\rho_i(w) = \rho_i$. The minimum function constitutes the required approximation to the *ground state energy density* $e(\rho_e, \rho_i)$. According to the thermodynamic stability conditions $e(\rho_e, \rho_i)$ should be a convex function of (ρ_e, ρ_i) . Since the functions $\rho_e \rightarrow e(w, \rho_e)$ are convex and the minimum of a set of convex functions is not necessarily convex, to guarantee the thermodynamic stability it is necessary to construct the convex envelope of the minimum function. It is clear that the choice of $\mathcal{W}_{\rho_i}$ is decisive for the quality of the approximation. We can choose $\mathcal{W}_{\rho_i}$ to be for instance the set $\mathcal{W}_{\rho_i, p}$ of all periodic ion configurations whose period does not exceed some p and $\rho_i(w) = \rho_i$. Then those points of the graph of the minimum function that coincide with the graph of the convex envelope can be associated with the periodic phases. The remaining points can be obtained as convex combinations of some other points belonging to the graph of the minimum function and therefore they are associated with mixtures of periodic phases. The canonical restricted phase diagrams of this kind have been obtained in [1] and [2]. It turns out however that the resulting canonical restricted phase diagrams are highly unstable. There are two main reasons for lack of stability in these diagrams. First, they do not allow for mixtures of periodic phases with different ion densities. In many instances the periodic phase corresponding to the point (ρ_e, ρ_i) of the restricted phase diagram can be shown to have higher energy than a suitable mixture of two or three periodic phases whose ion densities are different from ρ_i (cf. [3] and section 3). Second, the sets of admissible ion configurations used there are too restricted.

To overcome the stability problems mentioned and to avoid dealing with mixtures we have studied the system in the grand canonical ensemble and then translated the results into the canonical ensemble. In the *grand canonical ensemble* one is given the electron and ion chemical potentials (μ_e, μ_i) and the task is to find the minimum function of the set $\{f(w; \mu_e, \mu_i) : w \in \mathcal{W}\}$ of the free energy densities $f(w; \mu_e, \mu_i)$, where

$$f(w; \mu_e, \mu_i) = e^{(U)}(w; \mu_e) - \mu_e \rho_e(w; \mu_e) - \mu_i \rho_i(w) \quad (6)$$

and $\mathcal{W}$ is a set of admissible configurations. The thermodynamic stability requires the minimum function to be a concave function of (μ_e, μ_i) . Since the free energy $f(w; \mu_e, \mu_i)$ is a concave function of (μ_e, μ_i) and the minimum function of any set of concave functions is concave, the thermodynamic stability condition is satisfied automatically. We have taken for the class $\mathcal{W}$ all periodic ion configurations whose periods do not exceed 16 lattice constants, typically, however we have also constructed some diagrams where the maximal period has been increased to 40.

The canonical phase diagrams have a U -independent inversion *symmetry* about the point $(1/2, 1/2)$. It follows from (4) and (5) that, if $\tilde{w}$ is the ground state configuration of the ions at the point (ρ_e, ρ_i) , then the adjoint configuration $\tilde{w}^*$ is the ground state configuration at $(1 - \rho_e, 1 - \rho_i)$.

2 Some remarkable phases and their ions' configurations

To each point (ρ_e, ρ_i) the canonical phase diagram assigns a phase (more precisely a class of equivalent phases that consists of the phases related by the symmetries of the lattice), which is called here (ρ_e, ρ_i) -phase or the *stable phase*. Since we are concerned mainly with the ion subsystem, we characterize the phases by their ions' arrangements w and the electron density ρ_e . From a few thousand of the periodic ion configurations that have been tested, each time a restricted phase diagram was constructed, only a few per cent, at most, happen to be the ions' arrangements of the phases that appear in the phase

diagram. Many phases turn out to be the members of some infinite families of phases whose periodic ion configurations comply with peculiar rules and the ion and electron rational densities satisfy linear relations.

There are of course only two translationally invariant ion configurations, the *full configuration*, where all the sites of the lattice are occupied ($w(x) = 1$) and the *empty configuration* with no ions at all ($w(x) = 0$). The phases with translationally invariant ion configurations are called the *full phases* and the *empty phases*, respectively. The unique band of the empty configuration extends from -2 to 2 , while the band of the full configuration is obtained by translating the latter by $-U$. In the full or empty phases the system is a conductor, unless $\rho_e = 0$ or 1 . The spectrum of any ion configuration is confined to the interval given by the lower band edge of the full configuration, $-2 - U$, and the upper band edge of the empty configuration, 2 . For $U > 4$ the two bands do not overlap and the gap separating them opens in the spectrum of any other configuration.

From now on when we speak of periodic phases or periodic ion configurations we mean that their period is at least 2 .

The infinite families of the *ion n -molecule most homogeneous phases* $[p/q]_n$ are characterized by the so called *ion n -molecule most homogeneous configurations* of ions and the relation $\rho_i = n\rho_e$, where n is an integer, $\rho_e = p/q$ and $0 \leq \rho_e \leq 1/n$. In the context of the Falicov-Kimball model the ion n -molecule most homogeneous configurations have originally been defined in [12] (the case $n = 1$) and in [3] ($n \geq 2$). Let p and q , $p < q$, be relatively prime integers. The unit cell of the ion n -molecule most homogeneous configuration, whose ion density is $\rho_i = np/q$, has the length q and the arrangement of the ions is given by the solutions k_j to the equations

$$pk_j = j \bmod q, \quad j = 0, 1, \dots, np - 1. \quad (7)$$

The ions form clusters, the ion n -molecules, that consist of n consecutive sites occupied by the ions. The relation between the ion and electron densities means that in these phases there is one electron per ion n -molecule. The distances between any two nearest

neighbour clusters are either d or $d + 1$. Similarly the distances between next nearest neighbour n -molecules are either d' or $d' + 1$, and so on. In [3] the ion 1-molecule most homogeneous configurations have been called the *atomic most homogeneous configurations*. The following example illustrates the definitions introduced: for $p = 2$ and $q = 7$ the unit cell of $[2/7]_1$ phase is $[\bullet \circ \circ \circ \bullet \circ \circ]$, the unit cell of $[2/7]_2$ phase is $[\bullet \bullet \circ \circ \bullet \bullet \circ]$, and the unit cell of $[2/7]_3$ phase is $[\bullet \bullet \bullet \bullet \bullet \bullet \circ]$.

The infinite families of the *hole n -molecule most homogeneous phases* $[p/q]_n^*$ are characterized by the *hole n -molecule most homogeneous configurations* of the ions and the relation $1 - \rho_i = n\rho_e$, where n is an integer, $\rho_e = p/q$, $0 \leq \rho_e \leq 1/n$. The hole n -molecule most homogeneous configuration is defined in a similar way as the ion n -molecule most homogeneous configuration, except that the ions are replaced by the holes. In this sense the star can be interpreted as the operation of taking adjoint. Clearly the adjoint of the adjoint phase is the original phase. In these phases there is also one electron per molecule but this time it is the hole n -molecule.

In each family of the phases described so far the corresponding periodic ion arrangements can be viewed as composed of the same sort of molecules and there is one electron per molecule. There appear however in the phase diagrams other families of phases, where the corresponding periodic ion arrangements can be viewed as composed of the hole n -molecules, with $n \geq 2$, and of the composite molecules that consist of 1 ion, 1 hole and 1 electron. In these phases there is also one electron per each composite molecule but the number of electrons per hole molecule depends on its size. Specifically for a pair of irreducible rationals p/q and h/k , such that $1/3 \leq h/k \leq p/q \leq 1/2$ and $k - 2h = 1$, the $[p/q]_{h/k}$ phase has period q , the electron density $\rho_e = p/q$, the ion density $\rho_i = k\rho_e - h$, and the positions of the ions in its unit cell are given by the solutions k_j to the equations

$$p k_j = j \bmod q, \quad j = 0, 1, \dots, kp - hq - 1. \quad (8)$$

The rationals h/k of $[p/q]_{h/k}$ phases that are stable in our restricted diagrams belong to a special sequence, which will be specified later. The $[p/q]_{h/k}$ phase can be viewed as

a periodic arrangement of the hole k -molecules with h electrons per hole molecule and composite molecules built of 1 ion, 1 hole and 1 electron. This definition of $[p/q]_{h/k}$ families of phases does not apply to the limiting case $p/q = h/k = 1/2$ whose definition is postponed to the text that follows.

In the phase diagrams there appear also the families of phases that are adjoint to the $[p/q]_{h/k}$ phases. Every $[p/q]_{h/k}$ phase has its adjoint phase $[p/q]_{h/k}^*$, where the hole k -molecules are replaced by the ion k -molecules. The ion and electron densities of the $[p/q]_{h/k}^*$ phase satisfy the linear relation $\rho_i = -k\rho_e + k - h$.

One of the remarkable features of the families of periodic phases described so far is that their density pairs (ρ_e, ρ_i) and their ions' configurations can be obtained recursively, following a recursive construction of the so called Farey sequence. Therefore we can speak of the *Farey sequences of phases* like the Farey sequence of $[p/q]_n$ phases or the Farey sequence of $[p/q]_{h/k}$ phases, etc.

The *Farey sequence* $\mathcal{F}_n$ of order n is the ascending sequence of irreducible rational fractions between 0 and 1 whose denominators do not exceed n , i.e. the fraction p/q , with p and q relatively prime integers, belongs to $\mathcal{F}_n$ if $0 \leq p \leq q \leq n$ [13]. Having any two consecutive elements p/q and p'/q' of $\mathcal{F}_n$, the "parents", we can form their "descendant" p''/q'' ,

$$\frac{p''}{q''} = \frac{p + p'}{q + q'}, \quad (9)$$

which satisfies the inequalities

$$\frac{p}{q} < \frac{p''}{q''} < \frac{p'}{q'}. \quad (10)$$

This suggests how to construct $\mathcal{F}_n$ recursively. To build up $\mathcal{F}_n$ we start with $\mathcal{F}_1$, which is simply the pair of the "first parents" $\{0/1, 1/1\}$, and add to it the descendant $1/2$ of the "first parents", what results in $\mathcal{F}_2$. Then we choose a new pair of "parents" and repeat this step. The process is continued until the denominators of new "descendants" exceed n . Let p/q and p'/q' ($p/q < p'/q'$) be two consecutive elements of $\mathcal{F}_n$, for some n . Replacing the pair of the "first parents" $\{0/1, 1/1\}$ by the pair $\{p/q, p'/q'\}$ and continuing the recursive

process described until the denominators of new "descendants" exceed some $n' > n$, we end up with the *Farey sequence of order n' generated by the "parents" $\{p/q, p'/q'\}$* . The sequence so defined consists of those elements p''/q'' of $\mathcal{F}_{n'}$ that satisfy the inequalities $p/q \leq p''/q'' \leq p'/q'$. This definition can be extended to the vector case. Namely starting with the pair of vectors $\{(p/q, r), (p'/q', r')\}$ (with $p/q, p'/q'$ as above), where $0 \leq r, r' \leq 1$ are rationals, and forming "descendants" componentwise we arrive at the *vector Farey sequence of order n' ($n' > n$) generated by the pair of vector "parents"*. The recursion step is repeated until the denominators of the first components of the "descendants" exceed n' .

For each family of the phases described so far the density pairs (ρ_e, ρ_i) are of the form $(p_e/q_e, p_i/q_e)$, where p_e/q_e is an irreducible rational, and the components of these density pairs satisfy a linear relation $a\rho_e + b\rho_i = c$. Clearly, if (ρ_e, ρ_i) and (ρ'_e, ρ'_i) are two such pairs, then their "descendant" (ρ''_e, ρ''_i) (defined componentwise) satisfies the same linear relation. Therefore the density pairs of the families of phases considered constitute the vector Farey sequences. Specifically the density pairs of the $[p/q]_n$ phases form the vector Farey sequence generated by the "first parents" $\{(0/1, 0/1), (1/n, n/n)\}$, the density pairs of $[p/q]_n^*$ phases form the vector Farey sequence generated by $\{(0/1, 1/1), (1/n, 0/n)\}$, while those of the $[p/q]_{h/k}$ and $[p/q]_{h/k}^*$ phases form the vector Farey sequences generated by $\{(h/k, 0/k), (1/2, 1/2)\}$ and $\{(h/k, k/k), (1/2, 1/2)\}$, respectively, provided that h/k and $1/2$ are "parents" in $\mathcal{F}_k$.

It appears that following these recursive procedures one can construct also the unit cells of the phases considered. For that it is enough to specify the unit cells of the "first parents" and to identify the operation of forming descendants with the simple concatenating of these unit cells. In the case of $[p/q]_n$ phases the unit cell of the $(0/1, 0/1)$ phase is an empty site with no electrons $[o]$, while the unit cell of the $(1/n, n/n)$ phase consists of n consecutive occupied sites with one electron $[\overbrace{\bullet \bullet \dots \bullet}^n]$. Thus, for $n = 2$ their descendant has the densities $(1/3, 2/3)$ and its unit cell is $[o \bullet \bullet]$, the descendant

of $\{(0/1, 0/1), (1/3, 2/3)\}$ has the densities $(1/4, 1/2)$ and the unit cell $[\circ \circ \bullet\bullet]$ while the descendant of $\{(1/4, 2/4), (1/3, 2/3)\}$ has the densities $(2/7, 4/7)$ and the unit cell $[\circ \circ \bullet\bullet \circ \bullet\bullet]$, etc. In the case of $[p/q]_{h/k}$ phases the unit cell of $(h/k, 0/k)$ phase consists of k unoccupied sites with h electrons and that of $(1/2, 1/2)$ phase consists of two sites with one ion and one electron.

The $[p/q]_{h/k}$ phases seem to be natural generalizations of the n -molecule most homogeneous phases. They are generated just by different "first parents". Therefore it is tempting to name the phases generated by some pair of "first parents" phases by means of the recursive concatenating the *generalized most homogeneous phases*.

If on the other hand, given a pair of "first parents" phases, we first concatenate some number of the unit cells of one parent and then we concatenate the resulting composite cell with a similarly obtained composite cell of the other parent, we obtain a phase, which, compared to the corresponding most homogeneous phase, is extremely inhomogeneous. We term it *the generalized crenel phase*. The *crenel phases* considered in [3] are generated by the unit cells $\{[\circ], [\bullet]\}$. They are the "most inhomogeneous" phases among all the periodic phases with given density and period. It is clear that the families of the generalized most homogeneous phases and the generalized crenel phases are not disjoint.

Now we are in a position to define another family of the generalized most homogeneous phases, denoted $[p/q]_{1/2}$. All the members of this family have the same electron density $\rho_e = 1/2$ and therefore they are characterized by their ion density $\rho_i = p/q$. Using the recursive procedure of forming descendants, the density pairs of these phases can be obtained from the "first parents" $\{(1/2, 0/2), (1/2, 1/2)\}$ while their unit cells are generated by the "first parents" unit cells of the form $[\circ\circ]$ with 1 electron and $[\bullet\circ]$ with 1 electron. For instance the ion densities of the $[p/q]_{1/2}$ phases with $1/2 \leq p/q \leq 3/4$ and whose periods do not exceed 14 constitute the following sequence (in the ascending order): $1/2, 8/14, 7/12, 6/10, 5/8, 9/14, 4/6, 7/10, 10/14, 3/4$. The denominators of these fractions are equal to the periods of the corresponding phases. All these phases,

except those whose ion densities are $9/14$, $7/10$ and $10/14$, are simultaneously of the generalized crenel type.

The only *aperiodic phases* that appear in our restricted phase diagrams are those that constitute some mixtures of a few periodic or translationally invariant phases. The ion configurations of the mixtures are the corresponding mixtures of the ion configurations of the component phases [3]. A distinguished role is played by the mixtures of the translationally invariant phases, the so called *segregated phases* [1]. For $\rho_e < 1/2$ only the mixtures of the full phases and the vacuum turn out to be stable [3]. Their ion configurations are the corresponding mixtures of the full and empty configurations, i. e. all the ions clump together. It has been proved in [12] that if $\rho_e \neq \rho_i$ then there is a sufficiently large U such that above it the segregated phase is stable. In [3] the exact stability region of the segregated phases has been conjectured. This region consists of those points (ρ_e, ρ_i) that satisfy the inequality $\rho_e < \rho_i b(U)$, where $b(U)$ is the solution to the equation derived in [3]. More details concerning these phases can be found in [1], [3], [7], [12].

3 Main observations and conjectures

The excerpts from our data are presented in Figs 1–7. First, in Figs 1a and 1b, we show two schematic phase diagrams, which indicate where the families of characteristic phases, described in the previous section, are located in the (ρ_e, ρ_i) -plane.

Then in Figs 2–6 we show the actual data in the form of the restricted canonical phase diagrams. The set of the admissible ion configurations used to obtain these diagrams consisted of all the periodic ion configurations with the period not greater than 14 and of the full and empty configurations. In these diagrams the (ρ_e, ρ_i) -phases are represented by the black spots. In order to determine their rational ρ_e and ρ_i coordinates it is sufficient to know that the ρ_e values of the stable phases constitute the following sequence of irreducible rationals (in the ascending order):

$1/14$, $1/13$, $1/12$, $1/11$, $1/10$, $1/9$, $1/8$, $1/7$, $2/13$, $1/6$, $2/11$, $1/5$, $3/14$, $2/9$, $3/13$,

1/4, 3/11, 2/7, 3/10, 4/13, 1/3, 5/14, 4/11, 3/8, 5/13, 2/5, 5/12, 3/7, 4/9, 5/11, 6/13, 1/2. Moreover, if $\rho_e = p_e/q_e$ is any element of this sequence, then the ρ_i coordinates of the majority of the black spots, whose ρ_e coordinates are equal to p_e/q_e , belong to the sequence $1/q_e, 2/q_e, \dots, (q_e - 1)/q_e$. For the set of the admissible ion configurations used to obtain the diagrams presented the exceptions are rare (most of them are the phases with $\rho_e = 1/2$) and they are indicated in the captions of Figs 2–6. The periodic ion configurations corresponding to the (ρ_e, ρ_i) -phases with the ρ_e and ρ_i coordinates belonging to the sequences quoted can easily be determined. Suppose that $\rho_e = p_e/q_e$ and $\rho_i = p/q_e$. Then the unit cell has the length q_e . If we label the positions of the lattice sites in the unit cell $0, 1, \dots, p - 1$, then the positions of the ions are $0, k_1, \dots, k_{p-1}$, where k_1 is the inverse of $p_e \bmod q_e$ and $k_i = ik_1 \bmod q_e$ (cf the equations 12 and the comments of point 2 in the sequel).

Finally in Fig. 7 we show the intersections of the restricted canonical phase diagrams with the line $\rho_i = 1/2$, obtained for different U values.

1. The canonical phase diagram establishes a one to one U -dependent correspondence between the density pairs (ρ_e, ρ_i) and the stable phases. Suppose that at the point (ρ_e, ρ_i) , where $\rho_e = p_e/q_e$ and $\rho_i = p_i/q_i$ are irreducible rationals, the stable phase is periodic. The stable phase has necessarily the lowest Fermi sea energy. Now according to the second order degenerate perturbation theory with respect to U , for given $\rho_e = p_e/q_e$ the lowest Fermi sea energy among the periodic ion arrangements with $\rho_i = p_i/q_i$ is attained by the periodic configuration that has a gap at the Fermi level and this gap is the largest [14]. Specifically, to meet the requirement of the maximal gap, the period of the phase corresponding to $(p_e/q_e, p_i/q_i)$ has to be equal to Q – the least common multiple of q_e and q_i and the configuration w has to maximize $|\hat{w}(2\pi\rho_e)|$, where

$$\hat{w}\left(2\pi\frac{p_e}{q_e}\right) = \sum_{x=0}^{Q-1} \exp\left(i2\pi\frac{p_e}{q_e}x\right) w(x). \quad (11)$$

The same conclusion has been reached in [1]. If there is only one phase that complies with the rules quoted, we call it *uniform* (this is always the case if $Q = q_e$), if there is

more than one, we call them *degenerate uniform*. For all U values tested, from 0.2 up to 6, our restricted phase diagrams conform to these rules surprisingly well.

First, for all values of U , $0.2 \leq U \leq 6$, all the phases have appeared to be insulators. However, in order to reach this situation for small values of U the maximal period of the configurations tested has to be quite large. For instance for $U = 0.2$, the $(1/2, 1/2)$ -phase is stable also for ρ_e in a small vicinity of $1/2$, unless we take into account in the vicinity of $(1/2, 1/2)$ the phases whose periods go up to about forty. Since in the canonical phase diagrams presented in [1] the maximal period of the configurations tested is only 8, for small U the phases that appear are not insulators. In the diagrams presented in [2] this effect is even more pronounced, occurs also for U much larger than 1. This is a consequence of using too "directed" set of admissible ion configurations.

Second, only for intermediate U values, $1 < U < 3.6$ roughly, we have detected some *nonuniform* phases. One example is the $(3/7, 1/2)$ -phase whose ion configuration is not among those for which the maximum of $|\hat{w}(2\pi\rho_e)|$ is attained. It is visible for instance in $U = 3$ phase diagram, Fig.6. Its period 14 is the least common multiple of q_e and q_i , however its unit cell is of the form $[\bullet \circ \circ \bullet \circ \bullet \circ \circ \bullet \circ \bullet \bullet \bullet \circ]$, while the $(3/7, 1/2)$ phase that maximizes $|\hat{w}(2\pi\rho_e)|$ has the unit cell $[\bullet \circ \circ \bullet \circ \bullet \circ \bullet \circ \bullet \bullet \circ \bullet \circ]$. Other examples of nonuniform phases have been revealed by additional studies, not represented in the diagrams included. For U between 1 and 1.9 there appear $(1/2, \rho_i)$ -phases whose ion configurations are of the generalized crenel type, generated by the "parents" : $[\circ \bullet]$ with one electron and $[\bullet \bullet]$ (or $[\circ \circ]$) with one electron. The periods of these phases are not the least common multipliers of q_e and q_i .

2. For every $\rho_e = p_e/q_e$ there is a characteristic U value, $U_1(p_e/q_e)$, such that below $U_1(p_e/q_e)$ the general rule of point 1 is accompanied by an additional one. Namely, only the $(p_e/q_e, p_i/q_i)$ -phases such that $Q = q_e$ appear in the restricted phase diagrams. The smallest $U_1(p_e/q_e)$ is $U_1(1/2) \simeq 1$. As a consequence, below $U_1(p_e/q_e)$ the positions of the

ions in the unit cells of $(p_e/q_e, p_i/q_i)$ -phases are given by the solutions k_j to the equations

$$p_e k_j = j \bmod q_e, \quad j = 0, 1, \dots, p_i q_e / q_i - 1. \quad (12)$$

It is easy to see that once the k_1 solution has been found, the remaining solutions are given by $k_j = j k_1 \bmod q_e$. This additional rule turns out to be satisfied also for sufficiently large U (above $U = 3.6$ roughly), independently of ρ_e , simply because at these U values the ion arrangements of the periodic phases are the ion 1-molecule most homogeneous configurations.

Moreover, there is another characteristic U value, $U_2(p_e/q_e)$, $U_2(p_e/q_e) \leq U_1(p_e/q_e)$, such that below $U_2(p_e/q_e)$, if $p_{i,min}/q_e$ is the minimal ion density of the stable phases with $\rho_e = p_e/q_e$ and $p_{i,max}/q_e$ is the maximal one, then all the remaining $(p_e/q_e, p_i/q_e)$ phases with p_i assuming all the values $p_{i,min}, p_{i,min} + 1, \dots, p_{i,max}$ are stable as well. This is a more precise form of the observation made in [3]. The value of $U_2(p_e/q_e)$ decreases as the period q_e of the phases increases. It is not clear if there is some nonzero U_2 that is good for all periods. In Fig.3 ($U = 1.4$), for $\rho_e = 5/14$ there appear the phases corresponding to $p_i = 5, 6, 9, 10$ but those with $p_i = 7, 8$ are missing. Similarly, for $\rho_e = 4/11$ there appear the phases corresponding to $p_i = 4, 5, 7, 8$ but that with $p_i = 6$ is missing. Some other examples are visible in this diagram. All these missing phases are stable already in the phase diagram for $U = 0.8$, Fig.2. The values of $p_{i,min}$ and $p_{i,max}$ appear to be determined by the n -molecule most homogeneous phases and the generalized most homogeneous phases. For given $\rho_e = p_e/q_e$ the minimal $p_{i,min}/q_e$ and the maximal $p_{i,max}/q_e$ ion densities are determined by the minimal and maximal ρ_i values of the intersections of the line $\rho_e = p_e/q_e$ and the line segments formed by the $[p/q]_n$, $[p/q]_n^*$, $[p/q]_{h/k}$ and $[p/q]_{h/k}^*$ stable phases.

3. When the maximal period of the admissible ion configurations grows, then the number of the stable phases in the Farey sequences of the most homogeneous and the generalized most homogeneous phases grows as well. It appears that the order of any such a Farey sequence of phases is equal to the maximal period of the admissible

ion configurations. Moreover the Farey sequences of the most homogeneous and the generalized most homogeneous phases are stable in specific U -dependent intervals of ρ_e values. Only the ρ_e intervals of $[p/q]_1$ phases are nonvoid for all U . In the case of the ion n -molecule most homogeneous phases with $n > 1$ the critical U values at which the ρ_e intervals open have been determined in [3]. The hole n -molecule most homogeneous phases appear in the phase diagrams together with the 2-molecule most homogeneous phases, i.e. (according to [3]) below $U = 2.66$. The $[p/q]_n$ families appear in the phase diagrams in succession, starting with $n = 2$, so that for sufficiently small U only the phases with n not exceeding some characteristic value are stable [3]. In contrast, for given $U < 2.66$ all the $[p/q]_n^*$ phases with n greater than some characteristic value appear. For instance for $U = 2$ (Fig.4), those with $n = 1$ are still missing, but somewhere between $U = 2$ and $U = 1.4$ all the $[p/q]_n^*$ families become stable. The ρ_e stability intervals of the $[p/q]_n^*$ family of phases are determined by the points of intersection of the line $\rho_i = 1 - n\rho_e$ with the segments of those lines $\rho_i = n'\rho_e$ that correspond to the stable phases.

4. The Farey sequences of the $[p/q]_{h/k}$ generalized most homogeneous phases and their adjoint phases start to appear somewhere between $U = 1$ and $U = 0.8$, as U decreases. In Fig.2 ($U = 0.8$) three representatives of the $[p/q]_{1/3}$ phases and also three representatives of the adjoint family are visible. For $U = 0.4$ two families, $[p/q]_{1/3}$ and $[p/q]_{2/5}$ and their adjoints are well represented in our restricted phase diagrams and for $U = 0.2$ a few representatives of the $[p/q]_{3/7}$ family appear.

5. For U between 1 and 3.6 we have found a family of phases with $\rho_e = 1/2$. For $U > 2$ this family consists of the generalized most homogeneous phases of the $[p/q]_{1/2}$ type (and the adjoint family). For instance, in Fig.6 ($U = 3$) and in Fig.5 ($U = 2.6$) this family is represented by the phases whose ion densities (in the ascending order) are $8/14$, $7/12$, $6/10$, $5/8$, $9/14$ and $4/6$, where the denominators are equal to the periods of these phases. The phases that correspond to all ρ_i fractions quoted, except $9/14$ are uniform (the numerators of these ρ_i fractions exceed the halves of the denominators by

one) and their unit cells are of the generalized crenel type. In contrast the $(1/2, 9/14)$ -phase is degenerate uniform and it is not generalized crenel. Its unit cell has the form $[\circ \bullet \circ \bullet \bullet \bullet \circ \bullet \circ \bullet \bullet \bullet]$, in agreement with the recursive concatenating procedure. For sufficiently small U the degenerate uniform $[p/q]_{1/2}$ phases that are not generalized crenel transform into (degenerate uniform) phases of the generalized crenel type (with the same ion density). Thus in some points $(1/2, \rho_i)$ we observe the *structural phase transitions* – one stable periodic phase transforms into another stable periodic phase as U varies. For instance, for $U = 2.1$ the $(1/2, 9/14)$ -phase has the unit cell shown above. Already for $U = 1.9$ the unit cell of the $(1/2, 9/14)$ -phase is of the generalized crenel type: $[\circ \bullet \circ \bullet \circ \bullet \circ \bullet \bullet \bullet \bullet]$. Another manifestations of the effect described can be found in the diagrams for $U = 2$ (Fig.4) and for $U = 1.4$ (Fig.3). In Fig.4 the $(1/2, \rho_i)$ -phase with the largest ion density, $\rho_i = 7/10$, is degenerate uniform and generalized crenel. Its unit cell has the form $[\circ \bullet \circ \bullet \circ \bullet \bullet \bullet \bullet]$, while the unit cell of the generalized most homogeneous phase $[7/10]_{1/2}$ is $[\circ \bullet \circ \bullet \bullet \bullet \circ \bullet \bullet]$. The remaining $(1/2, \rho_i)$ -phases, with ρ_i equal to $5/8$, $6/10$, $7/12$ and $8/14$, are uniform and simultaneously generalized most homogeneous and generalized crenel. The only $(1/2, \rho_i)$ -phase visible in Fig.3, with $\rho_i = 10/14$, is also degenerate uniform and generalized crenel. Its unit cell is $[\circ \bullet \circ \bullet \circ \bullet \circ \bullet \bullet \bullet \bullet]$, while the $[10/14]_{1/2}$ -phase has the unit cell $[\circ \bullet \circ \bullet \circ \bullet \bullet \bullet \bullet \circ \bullet \bullet]$.

6. When U increases towards 4, all the periodic phases different from the 1-molecule most homogeneous ones start to disappear (their domains in the (μ_e, μ_i) plane shrink quickly). Thus, our diagrams suggest that above $U > 4$ the only phases that remain in the phase diagrams are the ion 1-molecule most homogeneous phases with arbitrary ρ_e , the mixtures of the ion 1-molecule most homogeneous phases and the full phases or the empty phases, and the segregated phases. This is in contrast with the canonical phase diagrams of [1] and [2].

7. The "evolution" of the phase diagrams with decreasing U can be viewed as building up an extra symmetry – the ion (or the electron) hole-particle symmetry. In the (ρ_e, ρ_i) -plane a polygon is formed, which is invariant with respect to the reflections in the line $\rho_i = 1/2$. The segments of the lines of the ion and hole most homogeneous phases determined by the mutual intersections of these lines constitute the sides of that polygon. For instance, in the $U = 1.4$ restricted phase diagram (Fig.3) the tetragon with the following vertices $(1/2, 1/2)$ $(1/3, 2/3)$ $(1/4, 2/4)$ is well visible, however it is not yet reflection invariant, since some phases inside this tetragon are missing, but for $U = 0.8$ (Fig.2) the reflection symmetry with respect to the line $\rho_i = 1/2$ is already restored. As U decreases the invariant polygon gradually extends through out the smaller ρ_e values. For $U = 0.4$ it consists of two tetragons whose boundaries are visible in the schematic phase diagram, Fig.1. The appearance of the hole most homogeneous phases $[p/q]_n^*$, the generalized most homogeneous phases $[p/q]_{h/k}$ and $[p/q]_{h/k}^*$ phases, as well as the fact that, as U decreases the minimal ρ_e values for which the $[p/q]_n$ phases are stable increase, can be viewed as manifestations of the tendency to restore the ion hole-particle symmetry. Because of the inversion symmetry of the phase diagram, the ion hole-particle symmetry is equivalent to the electron hole-particle symmetry.

8. In agreement with the Gibbs phase rule for a two component system at a fixed temperature, only mixtures of two or three phases appear in the restricted canonical phase diagrams. Actually in the diagrams presented, Figs 2–6, there appear also mixtures of more than three periodic phases. This is a consequence of the finite step with which the electron chemical potential axis has been scanned, when the grand canonical phase diagrams were constructed. The stable mixtures can be arranged in five groups. For $\rho_e < 1/2$ there appear mixtures of: (a) two periodic phases, (b) three periodic phases, (c) a periodic phase and a translationally invariant phase (where the electron density of the latter may vary in an interval), (d) two periodic phases with a translationally invariant

phase (whose electron density is uniquely determined by the periodic phases) and (e) the segregated phases that are mixtures of the vacuum (i.e. the empty phase with no electrons) and the full phases with ρ_e not greater than certain U -dependent critical value [3], [7], [12]. The mixtures belonging to the groups (a) and (b) are the insulators while those belonging to the other groups are the conductors. Typically the periodic phases that appear in the mixtures of the kind (c) and (d) belong to the Farey sequences of the (generalized) most homogeneous phases. When the maximal period of the admissible ion configurations grows, then the areas occupied by the mixtures of the kind (b) and (d) seem to shrink quickly. The mixtures of the kind (d), where the electrons densities of the two periodic phases are equal and assume a limiting value of the ρ_e stability interval of one of the Farey sequences of phases, seem to be the only exceptions. In Fig. 2 these are for instance the mixtures of $(1/6, 1/3)$, $(1/6, 1/2)$ and the full phases, the mixtures of $(2/5, 3/5)$, $(2/5, 4/5)$ and the full phases, etc. Therefore we expect that in the limit of the infinite maximal period there are no mixtures of the kind (b) and those of the kind (d) are relatively rare.

9. As U increases, then typically the periodic phases and their mixtures that are located at the boundary of the (ρ_e, ρ_i) region, where the periodic phases and their mixtures are stable, lose against mixtures of other periodic phases with a full or an empty phase. This results in an *insulator-conductor transitions*. In the limit $U \rightarrow \infty$ the neutral phases (with arbitrary ρ_e) remain the only insulators. However the insulator-conductor transitions can occur also when U decreases. As first observed in [3], as U decreases the n -molecular most homogeneous phases, with electron densities sufficiently close to 0 (or 1) lose their stability against mixtures of n' -molecular most homogeneous phases ($n' > n$) with the empty phases. In the case $n = 1$ this occurs below $U = 2/\sqrt{3}$ [3]. This effect is visible for instance in Fig.2 ($U = 0.8$). In the region of small ρ_e the representatives of the neutral phases terminate for larger ρ_e than this is the case for the ion n -molecular phases with $n = 2, 3$.

10. If we start with the point $(p_e/q_e, 0)$ and move it up along the line $\rho_e = \text{const}$, then typically it first passes through infinitely many mixtures of some empty phases (with smaller ρ_e) and some periodic ones (with larger ρ_e). After that the point enters an interval of ρ_i values, where only periodic phases and their mixtures are the stable phases. It is not clear if this interval contains only not more than $q_e - 1$ periodic phases with the period q_e and mixtures of two consecutive periodic phases of this kind. It may not be connected, there can be subintervals where mixtures of some periodic phases and full or empty phases are stable. Finally the point passes through infinitely many mixtures of some periodic phases (with smaller ρ_e) and some full phases (with larger ρ_e) until it reaches eventually the full phase (if $\rho_e > b(U)$, [3]) or the segregated phases (if $\rho_e < b(U)$). Similar scenario, but obtained by moving the point $(0, 1/2)$ along the line $\rho_i = 1/2$ and then repeating this procedure for many values of U , is presented in Fig. 7. This kind of the restricted canonical phase diagram can directly be compared with the diagrams shown in [1] and [2].

Acknowledgments

The authors acknowledge the financial support of the State Committee for Scientific Research (Poland) under grant 2 P302 147 04. We thank Christian Gruber and Daniel Ueltschi for interesting discussions.

References

- [1] J. K. Freericks and L. M. Falicov, Phys. Rev. B **41**, 2163 (1990)
- [2] P. Farkašovský and I. Bat'ko, J. Phys.:Condens. Matter **5**, 7143 (1993)
- [3] C. Gruber, D. Ueltschi and J. Jędrzejewski, J. Stat. Phys. **76**, 125 (1994)
- [4] J. Lach, R. Łyżwa and J. Jędrzejewski, Phys. Rev. B **48**, 10783 (1993); Acta Phys. Polonica A **84**, 327 (1993)
- [5] L. M. Falicov and J. C. Kimball, Phys. Rev. Lett. **22**, 997 (1969)
- [6] T. Kennedy and E. H. Lieb, Physica A **138**, 320 (1986)
- [7] U. Brandt, J. Low Temp. Physics **84**, 477 (1991)
- [8] D. I. Khomskii, in *Quantum theory of solids*, ed. I. M. Lifshits, Mir, Moscow 1982
- [9] U. Brandt and R. Schmidt, Z. Phys. B **63**, 45 (1986)
- [10] R. Łyżwa, Phys. Lett. A **164**, 323 (1992); Physica A **192**, 231(1993).
- [11] J. Jędrzejewski, J. Lach and R. Łyżwa, Phys. Lett. **134** A, 319 (1989). Physica A **154**, 529 (1989)
- [12] P. Lemberger, J. Phys. A **25**, 15 (1992)
- [13] G. H. Hardy and H. M. Wright, *Theory of Numbers*, Oxford Univ. Press, Oxford 1938
- [14] N. W. Ashcroft and N. D. Mermin, *Solid State Physics*, Holt, Rinehart and Winston, New York 1976

Figure Captions

Fig. 1 a) Schematic canonical phase diagram for $U = 0.4$. The shaded regions correspond to the segregated phases. The ion n -molecule most homogeneous phases are distributed along the continuous line segments while the hole n -molecule most homogeneous phases along the dashed line segments. The $[p/q]_{h/k}$ phases are lying along the dashed-dotted line segments.

b) Schematic canonical phase diagram for $U = 4$. The only periodic phases that appear in the diagram are the ion 1-molecule most homogeneous phases. The points representing these phases constitute the line segment that extends from $(0, 0)$ to $(1, 1)$.

Fig. 2 The restricted canonical phase diagram for $U = 0.8$. This is a part of the complete diagram, which is sufficient for reconstructing the remaining part by means of the inversion symmetry (cf section 1). The black spots represent the periodic phases (whose period is at least 2). Those black spots whose corresponding phases coexist in the grand canonical phase diagram are joined by the straight line segments. The points lying on such a segment represent the mixtures of the two periodic phases that correspond to the ends of the segment. The straight line segments partition the phase diagram into convex areas, most of which are triangles. These triangles correspond to the coexistence points of three periodic phases in the grand canonical phase diagrams. The points located inside these triangles represent mixtures of the three phases that correspond to the vertices of the triangles. Some segments are missing and therefore there are areas which according to this restricted diagram are mixtures of more than three phases. This is an artefact related with the precision of the diagram. All the (ρ_e, ρ_i) -phases conform to the equations 12.

Fig. 3 The restricted canonical phase diagram for $U = 1.4$. This is a part of the complete diagram, which is sufficient for reconstructing the remaining part by means of the

inversion symmetry. Only the $(1/2, 10/14)$ -phase does not conform to the equations 12. It does not belong also to the $[p/q]_{1/2}$ family. It is generalized crenel and its unit cell is $[\circ \bullet \circ \bullet \circ \bullet \circ \bullet \bullet \bullet \bullet]$. See the caption of Fig. 2 and section 3 for more explanations.

Fig. 4 The restricted canonical phase diagram for $U = 2$. This is a part of the complete diagram, which is sufficient for reconstructing the remaining part by means of the inversion symmetry. All the phases with $\rho_e = 1/2$, except that with the largest ion density $\rho_i = 7/10$, belong to the family of $[p/q]_{1/2}$ phases. Their ion densities (in the ascending order) are: $8/14, 7/12, 6/10, 5/8$. The unit cell of the $(1/2, 7/10)$ -phase is $[\circ \bullet \circ \bullet \circ \bullet \bullet \bullet \bullet]$. See the caption of Fig. 2 and section 3 for more explanations.

Fig. 5 The restricted canonical phase diagram for $U = 2.6$. See the caption of Fig. 2 and section 3 for more explanations. All the phases with $\rho_e = 1/2$ belong to the family of $[p/q]_{1/2}$ phases. Their ion densities (in the ascending order) are: $8/14, 7/12, 6/10, 5/8, 9/14, 4/6$.

Fig. 6 The restricted canonical phase diagram for $U = 3$. See the caption of Fig. 2 and section 3 for more explanations. All the phases with $\rho_e = 1/2$ belong to the family of $[p/q]_{1/2}$ phases. Their ion densities (in the ascending order) are: $8/14, 7/12, 6/10, 5/8, 9/14, 4/6$. The $(3/7, 7/14)$ -phase is not uniform.

Fig. 7 This is a schematic phase diagram, the intersections of the restricted canonical phase diagrams with the line $\rho_i = 1/2$, obtained for the specific U values marked on the horizontal axis. We can distinguish a few kinds of stable phases. In the white region marked "segregated", the segregated phases are the only stable phases. In the light grey region marked "most homogeneous&full" a stable phase is a mixture of one or two ion n -molecule most homogeneous phases and a full phase. In the dark grey region marked "periodic phases and their mixtures" the stable phases are either periodic or mixtures of one or two periodic phases. Finally in the unmarked

grey region the stable phases can be either like in the dark or light grey regions, or something else, depending on ρ_e . For the specific U values marked on the horizontal axis, the critical ρ_e values, where the kind of stable phases changes, are marked by the short horizontal segments. The continuous segments correspond to periodic phases while the dotted ones to mixtures.